New Jersey Division of HIV/AIDS Services Persons Living with HIV/AIDS as of June 30, 2004 Bloomfield Residents

l. Current HIV/AIDS Category	Cases	elescent *		Pediat Cases	용			Tota ases	용
AIDS-Any OI AIDS-CD4 Only HIV Only	26 # #	19 # #		O # #	 (‡) #		26 40 70	19 29 51
Total	135	100		#		- #	_	136	100
Current Age * Cases %		[/] Eth.		Adult/ Cases	s %	Pediat	용	Tota Cases	al %
<5 # # 5-12 0 0 13-19 # # 20-29 6 4	White Black Hispa Other	e, Not Hisp k, Not Hisp	anic	54 # 20 #	40 # 15 #	O # O #	0 # 0 #	54 58 20 #	40 43 15 #
30-39 33 24 40-49 55 40 50+ 39 29 Unk. 0 0 Total 136 100	Total			135		#	#	136	100
. Patient Group		Ac Males		dolescer	nt Tran Female		ion M	odes ** Tota	
Men sex with men (MSM) Injection drug user (IDU) MSM/IDU Heterosexual contact Risk not reported/other***		37 16 # # 26	41 18 # # 29		12 (14	0 0 2 28 0 0 # # 4 33		3 2	 37 28 # #
Total		91	100		43	3 100		13	34 1
		Males	용	diatric	Female	es %		Tota	
Parent at risk/has AIDS/HIV Risk not reported/other***		# O	#		‡	# # D 0			# O
Total		 #	 #			 # #			 #

NOTE: Includes only those persons known to be infected with HIV.

Total

- # Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.
- Classification based on current age at the end of June 2004.
- ** 1 patients are currently of adult/adolescent age (13+ years old), but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1,2, and 3, and as pediatric reports in table 4.

#

#

 $\ensuremath{^{\star\star\star}}$ Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services AIDS Cases Reported as of June 30, 2004 Bloomfield Residents

Adult/Adolescent*

1. Disease C	ategory	Cases	(응)	Deaths (%)	Cases (%)	Deaths (%) Cases (응)	Deaths (%)
PCP Other OI KS Alone No Diseas		 	0 (29) # (#) # (#) 0 (35)	43 (88) # (#) # (#) 20 (36)	O (O) # (#) # (#) O (O)	0 (0 (0 (0 (0) # (#) #)	43 (88) # (#) # (#) 20 (36)
Total		171	1 (100)	106 (64)	# (100)	0 (0) 172 (1	00)	106 (63)
2. Age *	Cases	ଚ	3. Race	/Eth.	Adult/Adl* Cases %	Pediatric* Cases %	Total Cases	90	
Under 5 5-12 13-19 20-29 30-39 40-49 Over 49 Unknown	0 # 18 73 57 22 0	42 33	Blac Hisp Othe	r/Unknown 	# # # # 24 14 0 0 171 100	0 (# 96 0 52 0 24 0 0 #	56 30 14 0 	

Pediatric*

Total

4.	Patient	Group
----	---------	-------

Total

172 100

. Patient Group	Adult/Adolescent Transmission Modes **									
	Males	용	Females %	Total	용					
Men sex with men (MSM)	47	38	0 0	47	27					
Injection drug user (IDU)	38	30	19 41	57	33					
MSM/IDU	8	6	0 0	8	5					
Heterosexual contact	14	11	18 39	32	19					
Risk not reported/other***	18	14	9 20	27	16					
Total	125	100	46 100	171	100					

	Pediatric Transmission Modes						
	Males	용	Females %	Total	%		
Parent at risk/has AIDS/HIV	#	#	# #	#	#		
Risk not reported/other***	0		0 0	0	0		
Total	#	#	# #	#	#		

[#] indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.

Classification at time of AIDS diagnosis.

^{** 0} patients were diagnosed with $\widetilde{\text{AIDS}}$ as adults, but have evidence either of having been HIV-infected or of having acquired AIDS as children. They are counted as adult/adolescent reports in tables 1,2, and 3; and as pediatric reports in table 4.

^{***} Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services AIDS Cases Reported as of June 30, 2004 Bloomfield Residents

5. AIDS Cases by Year of Diagnosis.

Year of Diagnosis	Number of Cases	Number of Deaths	Case-Fatality Rate
Before 1985	#	#	#
1985	#	#	#
1986	#	#	#
1987	6	6	100%
1988	8	8	100%
1989	13	12	92%
1990	7	5	71%
1991	16	15	94%
1992	13	12	92%
1993	20	14	70%
1994	14	8	57%
1995	8	#	#
1996	13	#	#
1997	9	#	#
1998	#	#	#
1999	8	#	#
2000	8	#	#
2001	7	#	#
2002	#	#	#
2003	5	#	#
2004	#	#	#
Total	172	106	62%

New Jersey Division of HIV/AIDS Services HIV INFECTION-not-AIDS Reports with Identifiers Reported as of June 30, 2004 Bloomfield Residents

1. Age *	Cases	%	2. Race/Ethnicity	Adult/A		Pediat Cases	ric* %	Total Cases	%
Under 5	#	#	White, Not Hispanic	30	38	0	0	30	38
5-12	0	0	Black, Not Hispanic	#	#	#	#	34	44
13-19	#	#	Hispanic	11	14	0	0	11	14
20-29	15	19	Other/Unknown	#	#	#	#	#	#
30-39	26	33							
40-49	23	29	Total	78	100	#	#	79	100
Over 49	13	16							
Unknown	0	0							
Total	79	100							
3. Patient G	roup		Adul	Lt/Adoles	scent	Transmissi	on Mod	es **	

Risk not reported/other***	0 	π • 	0 	0 	υ Ο 	0
Parent at risk/has AIDS/HIV		#	 #	 #	 #	#
	Males	Pedia %	tric Transmi Females		les Total	olo
Total	51	100	27	100	78	100
Risk not reported/other***	19	37 	9	33	28	36
Heterosexual contact	#	#	#	#	#	#
MSM/IDU	#	#	0	0	#	#
Injection drug user (IDU)	7	14	9	33	16	21
Men sex with men (MSM)	19	37	0	0	19	24
	Males	% 	Females	% 	Total	%
Patient Group	Ac		escent Transı		lodes **	

- * Classification at time of first HIV+ test or at time of report if not yet confirmed HIV+.
- # indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.
- ** 0 patients were diagnosed with HIV as adults, but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1 and 2; and as pediatric reports in table 3.
- *** Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

NOTE: These tables include those reports of HIV Positive infection that have not yet been updated to AIDS status. Not included in these tables are cases initially reported as HIV-not-AIDS that have subsequently been updated to AIDS. Also not included in these tables are HIV Perinatal Exposure Reports where the serostatus of the child is not confirmed HIV Positive; i.e., the HIV serostatus of the child is undetermined or has been found to be negative (seroreverted).

New Jersey Division of HIV/AIDS Services HIV/AIDS Cases Reported as of June 30, 2004 Bloomfield Residents

1.	Current HI	V/AIDS Ca	tegory		Adult/Adol Cases	%		Ped Case	S	%	Cases			
	AIDS-Any O AIDS-CD4 O HIV Only	oI Only			# 60 #				# O #	# O #	112 60 79	! 4) 2	15 24 31	
	Total					100			#	#		. 10		
2.				3.	Race/Eth.		Cas	ses	용	Pediat Cases	용		ases	tal %
	Under 5 5-12 13-19 20-29 30-39 40-49 Over 49 Unknown	# 43 101 71 32 0	# # 17 40 28 13		White, Not Black, Not Hispanic Other/Unkn	Hispanic Hispanic own		# # 	# # # #	# # 			126 86 35 # 251	34 14 #
	Total	251												
4.	Patient Gr	-				Ad Males	용		Fe	Transmissi males %		Total	%	
	Men sex wi Injection MSM/IDU Heterosexu Risk not r	drug user	ISM) (IDU) tt			66 45 10 18 37	38 26 6 10 21			0 0 28 38 0 0 27 37 18 25			27 29 4 18	
						Males	Peo %	diatr	Fe	ansmission		Total		
	Parent at Risk not r		AIDS/HI			# O	#			# # 0 0		# O	# 0	

NOTE: Includes only those persons known to be infected with HIV.

- # Indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.
- * Classification at time of first HIV+/AIDS diagnosis.

Total

** 0 patients were diagnosed with HIV/AIDS as adults, but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1,2 and 3; and as pediatric reports in table 4.

#

 $\star\star\star$ Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services HIV/AIDS Cases Reported as of June 30, 2004 Bloomfield Residents

5. HIV/AIDS Cases by Year of First HIV/AIDS Diagnosis.

Year of First HIV/AIDS Diagnosis	Cases	Number Known Dead	Case-Fatality Rate
Before 1985	5	#	#
1985	5	5	100%
1986	9	8	89%
1987	12	9	75%
1988	10	10	100%
1989	20	19	95%
1990	14	8	57%
1991	14	10	71%
1992	22	14	64%
1993	18	8	44%
1994	14	#	#
1995	14	#	#
1996	12	#	#
1997	14	#	#
1998	10	#	#
1999	9	#	#
2000	9	#	#
2001	13	#	#
2002	13	0	0%
2003	11	#	#
2004	#	#	#
Total	251	115	46%